


# Usefulness of low-dose combined spinal epidural anesthesia for a woman with systemic lupus erythematosus, mechanical prosthesis in the mitral valve position, and worsening heart failure presenting for cesarean delivery

Devon Weidner, BS<sup>a</sup>, David Hepburn, DO<sup>b</sup>, Blake Harris, DO<sup>b</sup>, Jennifer Chae-Kim, MD<sup>c</sup>, Jessica Ehrig, MD<sup>d</sup>, and Michael P. Hofkamp, MD<sup>b</sup> 

<sup>a</sup>College of Medicine, Texas A&M University Health Science Center, Bryan, Texas; <sup>b</sup>Department of Anesthesiology, Baylor Scott & White Medical Center – Temple, Temple, Texas; <sup>c</sup>Department of Obstetrics and Gynecology, Baylor Scott & White Medical Center – Temple, Temple, Texas; <sup>d</sup>Division of Maternal Fetal Medicine, Department of Obstetrics and Gynecology, Baylor Scott & White Medical Center – Temple, Temple, Texas

## ABSTRACT

A 31-year-old woman, gravida one, para zero, at 32 weeks, 4 days gestation, with a history of antiphospholipid antibody syndrome, mitral valve replacement requiring anticoagulation, chronic diastolic heart failure, and systemic lupus erythematosus was admitted to the hospital for worsening cardiac decompensation with superimposed pneumonia. She was on warfarin for anticoagulation at the time of hospital admission and eventually started on an intravenous heparin infusion. Cesarean delivery was planned due to comorbidities and anticoagulation status. After administration of betamethasone for fetal lung maturity, the patient's heparin infusion was discontinued approximately 16 hours prior to cesarean delivery. Upon obtaining laboratory testing that confirmed appropriate coagulation status, a low-dose combined spinal epidural anesthetic technique was used for cesarean delivery and the expected hemodynamic shifts due to spinal anesthesia were mitigated with a prophylactic norepinephrine infusion.

**KEYWORDS** Analgesia; anesthesia; conduction; anesthesia; obstetrical; cesarean section

Neuraxial anesthesia is the recommended technique for cesarean delivery<sup>1</sup> but carries the risks of hypotension, infection, and, rarely, spinal hematoma.<sup>2</sup> Patients with preexisting cardiac conditions are susceptible to the deleterious effects of sympathectomy arising from spinal anesthesia, while patients with coagulopathy have an increased risk of spinal hematoma with spinal anesthesia. We present a patient with a history of systemic lupus erythematosus, lupus anticoagulant, and mechanical prosthesis in the mitral valve position with worsening heart failure and pneumonia who presented for cesarean delivery.

## CASE DESCRIPTION

A 31-year-old woman, gravida one, para zero, at 32 weeks, 4 days gestation, with a body mass index of 26.7 kg/m<sup>2</sup>, was admitted to the hospital for worsening cardiac decompensation with superimposed pneumonia. Her medical history was notable for systemic lupus erythematosus, antiphospholipid antibody syndrome, mitral valve replacement with a mechanical valve requiring continuous systemic anticoagulation with warfarin, and chronic diastolic congestive heart failure (New York Heart Association class II). An echocardiogram demonstrated normal left ventricular cavity size and mild global hypokinesis with a left ventricular

**Corresponding author:** Michael P. Hofkamp, MD, Department of Anesthesiology, Baylor Scott & White Medical Center – Temple, 2401 South 31st Street, Temple, TX 76508 (e-mail: [Rfums2004@hotmail.com](mailto:Rfums2004@hotmail.com))

The authors report no conflicts of interest. The patient provided written permission for this case report to be published.

Received April 24, 2021; Revised May 10, 2021; Accepted May 13, 2021.

ejection fraction of 46%. The patient had a SARS-CoV-2 test that did not show infection, and her pneumonia was empirically treated with azithromycin and ceftriaxone. She received a 2-day course of betamethasone to facilitate fetal lung maturity, in anticipation of preterm cesarean delivery. She was transitioned from warfarin to a heparin infusion to maintain therapeutic anticoagulation, and the infusion was discontinued approximately 16 hours prior to the scheduled cesarean delivery. On the morning of surgery, coagulation studies demonstrated an elevated activated partial thromboplastin time of 73.8 seconds but a normal anti-Factor Xa level of <0.1 IU/mL. The hematology service attributed the elevated activated partial thromboplastin time to the patient's prior history of having lupus anticoagulant and made the assessment that surgery and neuraxial anesthesia could be safely conducted due to the normal anti-Factor Xa level.

The patient had a scheduled elective cesarean delivery, and combined spinal epidural anesthesia was performed. Hyperbaric bupivacaine 7.5 mg was administered into the intrathecal space followed by placement of an epidural catheter, and a continuous norepinephrine infusion was started at 0.02 mcg/kg/min. The initial neuraxial dermatome level was T9 when checked by pinprick. Lidocaine 2% was administered through the epidural catheter in two sequential 5 mL aliquots, and a T4 dermatome level was eventually obtained. The patient had an uncomplicated intraoperative course and was discharged home on the fourth postoperative day.

## DISCUSSION

To our knowledge, this is the first report describing a parturient with systemic lupus erythematosus and mechanical prosthesis in the mitral valve position who presented for cesarean delivery. Anti-phospholipid syndrome is a systemic autoimmune disease characterized by arterial or venous thrombosis mediated by antibodies to phospholipids<sup>3</sup> and results in a hypercoagulable state.<sup>4</sup> Previous accounts of pregnant patients with mechanical heart valves reported a 43% complication rate that included valve thrombosis and systemic anticoagulation that resulted in uterine hemorrhage.<sup>5,6</sup> To balance the risk of thrombus with the risk of bleeding, a heparin infusion was used during the hospitalization to bridge the patient to cesarean delivery. The American Society of Regional Anesthesia recommends discontinuation of a heparin infusion 4 to 6 hours prior to a neuraxial procedure and confirmation of normal coagulation lab values,<sup>7</sup> and we chose to discontinue the infusion 16 hours before the surgical procedure to ensure that the patient would meet criteria for surgery. On the day of surgery, there was discordance between the activated partial thromboplastin time and the anti-Factor Xa level. The consultant hematologist made the assessment that the anti-Factor Xa was a more accurate

indicator of the patient's coagulation status<sup>8</sup> and that the elevated activated partial thromboplastin time was artifactually elevated due to a preexisting lupus anticoagulant and did not recommend thromboelastography.

Single-shot spinal anesthesia can cause an abrupt decrease in systemic blood pressure due to rapid blockade of sympathetic nerve fibers arising from thoracic spinal nerve roots.<sup>9</sup> To balance the competing goals of achieving a dense surgical block and maintaining hemodynamic stability, we decided to perform a low-dose combined spinal epidural anesthetic.<sup>10</sup> We were able to employ a low dose of intrathecal bupivacaine to provide a reliable surgical block with limited hypotension and we extended the block with incremental administration of lidocaine 2% through the epidural catheter.

## ORCID

Michael P. Hofkamp  <http://orcid.org/0000-0002-5749-2849>

1. Practice Guidelines for Obstetric Anesthesia: an updated report by the American Society of Anesthesiologists Task Force on Obstetric Anesthesia and the Society for Obstetric Anesthesia and Perinatology. *Anesthesiology*. 2016;124(2):270–300. doi:10.1097/ALN.0000000000000935.
2. Lagerkranser M. Neuraxial blocks and spinal haematoma: review of 166 case reports published 1994–2015. Part 1: demographics and risk-factors. *Scand J Pain*. 2017;15:118–129. doi:10.1016/j.sjpain.2016.11.008.
3. Chaturvedi S, McCrae KR. Diagnosis and management of the antiphospholipid syndrome. *Blood Rev*. 2017;31(6):406–417. doi:10.1016/j.blre.2017.07.006.
4. Di Prima FA, Valenti O, Hyseni E, et al. Antiphospholipid syndrome during pregnancy: the state of the art. *J Prenat Med*. 2011;5(2):41–53.
5. van Hagen IM, Roos-Hesselink JW, Ruys TP, et al. Pregnancy in women with a mechanical heart valve: data of the European Society of Cardiology Registry of Pregnancy and Cardiac Disease (ROPAC). *Circulation*. 2015;132(2):132–142. doi:10.1161/CIRCULATIONAHA.115.015242.
6. Gupta R, Ranchal P, Harburger J. Mechanical valve thrombosis in a pregnant patient: a case of therapeutic failure. *Cureus*. 2019;11:e5615.
7. Horlocker TT, Vandermeulen E, Kopp SL, Gogarten W, Leffert LR, Benzon HT. Regional anesthesia in the patient receiving antithrombotic or thrombolytic therapy: American Society of Regional Anesthesia and Pain Medicine evidence-based guidelines (fourth edition). *Reg Anesth Pain Med*. 2018;43(3):263–309. doi:10.1097/AAP.0000000000000763.
8. Chandramouli NB, Rodgers GM. Management of thrombosis in women with antiphospholipid syndrome. *Clin Obstet Gynecol*. 2001;44(1):36–47. doi:10.1097/00003081-200103000-00006.
9. Šklebar I, Bujas T, Habek D. Spinal anaesthesia-induced hypotension in obstetrics: prevention and therapy. *Acta Clin Croat*. 2019;58(Suppl 1):90–95. doi:10.20471/acc.2019.58.s1.13.
10. Van de Velde M. Low-dose spinal anesthesia for cesarean section to prevent spinal-induced hypotension. *Curr Opin Anaesthesiol*. 2019;32(3):268–270. doi:10.1097/ACO.0000000000000712.